

Forester

An emulsion in water formulation containing 100 g /l (10% w/w) cypermethrin for control of Large Pine Weevil (*Hylobius abietis*) in forestry



Hylobius, the Large Pine Weevil damages forestry transplants by feeding on the bark, often ring-barking the tree. Adult weevils have two peak periods of feeding, the first between April and June and the second between July and October.



Forester is a contact and stomach acting insecticide with residual activity, applied as a post planting treatment for forestry transplants.

Rate and method of application:

Post planting forestry treatment: 20ml product/litre knapsack or forest drench gun
(delivering 10-20 mls of solution to the base of the tree stem)

Timing of application

Post planting forestry applications should be made either before damage is seen or on the onset of damage in the first two years after transplanting. A maximum of three applications can be made in any one year. Applications are recommended in April and July.

Note: Forester has been tested on Sitka Spruce but safety to other tree species cannot be assumed. Test a small number of trees for tolerance first before treating large numbers of species of trees.

Contents: 1 and 5 Litres



HARMFUL

HARMFUL: if swallowed.

IRRITATING to skin.

MAY CAUSE SENSITISATION by skin contact.

VERY TOXIC TO AQUATIC ORGANISMS, may cause long-term adverse effects in the aquatic environment.

To avoid risks to man and the environment, comply with the instructions for use.



DANGEROUS FOR THE ENVIRONMENT

MAPP NO: 13164

Distributor:

Fargro Limited,
Toddington Lane,
Littlehampton,
West Sussex,
BN17 7PP
Tel: (01903) 721591
Fax: (01903) 730737
Email: promos@fargro.co.uk
Web: www.fargro.co.uk

Approval holder

Agriphar S.A.
Rue de Renory 26,
B-4102 Ougree
Belgium