

ACARICIDE



BORNEO[®]

A suspension concentrate containing 110g/L etoxazole.

For use as an acaricide for the control of spider mites
in protected tomatoes and aubergines.

Product registration number: **MAPP 13919**

New product in 2009.

Based on a new active ingredient to the UK.

Control of spider mites.

Effective on eggs, nymphs and larvae. Adult females become sterile.

Long lasting.

Contact and translaminar activity.

fargro[®]

Fargro Ltd
Toddington Lane
Littlehampton
West Sussex
BN17 7PP
Tel: 01903 721591
Fax: 01903 730737
www.fargro.co.uk

ALWAYS READ THE LABEL. USE PESTICIDES SAFELY

BORNEO

- First acaricide in UK in the diphenyloxazoline group.
- A chitin synthesis inhibitor - IRAC* group 10B. Acts on chitin synthesis so mites cannot moult. Eggs do not hatch.
- Activity believed to be fairly specific to mites.
- Effective on mite eggs, nymphs and larvae. Adults become sterile.
- Contact and translaminar activity.
- Long residual activity.
- Usable with a range of beneficial insects.

The following text is taken from the approved product label:

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

BORNEO is a contact acaricide with contact action for the control of red spider mite in protected tomatoes and aubergines. It inhibits the development of eggs and larval stages of mites (*Tetranychus urticae*) and can be used on indoor tomatoes and aubergines (egg plants).

The product will give best performance if used at the first appearance of the first mobile stages of spider mites.

BORNEO has residual action of around 45 to 60 days.

MIXING AND SPRAYING

Invert bottle before use to ensure a uniform mixture. Half fill spray tank with clean water, commence agitation and add the required amount of BORNEO. Maintain agitation during completion of filling and during application.

Rinse empty container thoroughly at least three times. Add washings to the spray

tank at time of filling. Do not leave spray liquid in the sprayer for long periods of time overnight or during meal breaks.

In case of a high adult mite population, an adult acaricide should be applied.

RATE OF USE

Tomatoes and Aubergines

BORNEO should be used at a concentration of 25-30 ml per 100 L of water, in 500-1500 L water/ha; or at a concentration of 35 ml per 100 L of water in 500-1000 L water/ha. The higher concentration of 35ml/100L water should be applied when a high pest population is present.

Use sufficient water volume to obtain complete crop coverage.

RESISTANCE MANAGEMENT

In order to avoid the development of resistance, it is recommended that BORNEO is used only once per year

IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL ACARICIDE

CROPS: Tomatoes and Aubergines grown as protected crops.

Maximum Individual Dose : 35 ml per 100 litres of water (see other specific restrictions).

Other Specific Restrictions: A maximum individual dose of 500ml/ha must not be exceeded.

Maximum number of applications : 1 per crop.

Harvest Interval : 3 days.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE GLOVES when handling the concentrate or when applying the product. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH HANDS before meals and after work.

Keep locked up and out of the reach of children.

Keep away from food, drink and animal feeding stuffs.

When using do not eat, drink or smoke.

Environmental Protection

Do not contaminate water with the product or its container.

Storage and Disposal


KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

DO NOT REUSE CONTAINER FOR ANY OTHER PURPOSE.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Borneo A suspension concentrate containing 110g/L etoxazole.



**DANGEROUS
FOR THE
ENVIRONMENT**

**VERY TOXIC TO AQUATIC ORGANISMS, MAY CAUSE
LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVI-
RONMENT.**

This material and its container must be disposed of as haz-
ardous waste. Avoid release to the environment.
Refer to special instructions / safety data sheets.

To avoid risks to man and the environment, comply with the instructions for use.

CONDITIONS OF SUPPLY

All good supplied by us are of high grade and we believe them to be suitable but as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether they supervise or assist in the use of such goods.

®BORNEO is a registered trademark of
Sumitomo Chemical Co Ltd.

Approval Holder:

Sumitomo Chemical Agro Europe S.A.S.
Parc d'affaires de Crécy
2 rue Claude Chappe
69370 St Didier au Mont d'Or
France.

Pack Size: 250 ml
Outers of: 12

Marketed in the UK by :
Interfarm UK Ltd
36, Newgate Street
Doddington, Cambs PE15 0SR
Tel 01354 741414

*IRAC Insecticides Resistance Action
Committee

Safety data sheet available on request.

Distributed by Fargo Ltd.

© 2009 Copyright of Fargo and its suppliers.